

Joint law of the hitting time, overshoot and undershoot for a Lévy process

Laure Coutin*, Waly Ngom†

March 9, 2016

Abstract

Let $(X_t, t \geq 0)$ be a Lévy process which is the sum of a Brownian motion with drift and a compound Poisson process. We consider the first passage time τ_x at a fixed level $x > 0$ by $(X_t, t \geq 0)$, and $K_x := X_{\tau_x} - x$ the overshoot and $L_x := x - X_{\tau_x-}$ the undershoot. We first study the continuity of the density of τ_x . Secondly, we calculate the joint law of (τ_x, K_x, L_x) .

keywords: Lévy process, jump process, hitting time, overshoot, undershoot.

1 Introduction

In the theory of risk in continuous time the surplus of an insurance company is modelled by a stochastic process $(X_t, t \geq 0)$. The positive real number x denotes the initial surplus and $\tau_x := \inf\{t \geq 0 : X_t \geq x\}$ may be interpreted as the default time. This paper deals with τ_x when X is a Lévy process, sum of a drifted Brownian motion and a compound Poisson process. Our main results lead to the regularity of the density of the hitting time and to an explicit expression characterizing the joint distribution of the triplet (first hitting time, overshoot, undershoot).

J. Bertoin [1] gives a quick and concise treatment of the core theory on Lévy processes with the minimum of technical requirements. He gives some details on subordinators, fluctuation theory, Lévy processes with no positive jumps and stable processes.

P. Tankov and R. Cont [4] provide a self-contained overview of theoretical, numerical and empirical research on the use of Lévy processes in financial modeling. When the process X has jumps, the first results are obtained by Zolotarev [17] and Borovkov [3] for X a spectrally negative Lévy process. Moreover, if X_t the probability density with respect to the Lebesgue measure $p(x, t)$ then the law of τ_x has the density with respect to the Lebesgue measure $f(t, x)$ such that

*IMT, University of Toulouse, France, laure.coutin@math.univ-toulouse.fr

†FST, University Cheikh Anta Diop of Dakar, Sénégal, ngomwaly@gmail.com- IMT, University of Toulouse, France, waly.ngom@math.univ-toulouse.fr

$xf(t, x) = tp(x, t)$. R. A. Doney [6] deals with hitting probabilities, hitting time distributions and associated quantities for Lévy processes which have only positive jumps. He gives an explicit formula for the joint Laplace transform of the hitting time τ_x and the overshoot $X_{\tau_x} - x$.

When X is a stable Lévy process, Peskir [14] obtains an explicit formula for the passage time density. Moreover, if X has no negative jumps and if $S_t = \sup_{0 \leq s \leq t} X_s$ is its running supremum, Bernyk et al. [2] show that the density function f_t of S_t can be characterized as the unique solution to a weakly singular Volterra integral equation of the first kind.

In the case where X is a jump-diffusion process, with jump size following a double exponential law, Kou and Wang [11] give the law of τ_x . They obtain explicit solutions of the Laplace transform of the distribution of the first passage time. Laplace transform of the joint distribution of jump-diffusion and its running maximum, $S_t = \sup_{s \leq t} X_s$, is also obtained. Finally, they give numerical examples.

For a general Lévy process, Doney and Kyprianou [7] and Kyprianou [13] give the law of the quintuplet $(\bar{G}_{\tau_x}, \tau_x - \bar{G}_{\tau_{x-}}, X_{\tau_x} - x, x - X_{\tau_{x-}}, x - \bar{X}_{\tau_{x-}})$ where $\bar{X}_t = \sup_{s \leq t} X_s$ and $\bar{G}_t = \sup\{s < t, X_s = X_t\}$.

For a stable Lévy process X of index $\alpha \in (1, 2)$ the Lévy measure of which has the density $s(x) = cx^{-\alpha-1}$, $x > 0$, R. A. Doney in [8] considers the supremum $S_t = \sup_{s \leq t} X_s$ of X . He shows that S_1 behaves as $s(x) \sim cx^{-\alpha-1}$ as $x \rightarrow +\infty$.

Recently, Pogany, Tibor K and Nadarajah in [15] give a shorter and more general proof of R. A. Doney's previous result [8]. They derive the first known closed form expression for $s(x)$ and the corresponding cumulative function, then they obtain the order of the remainder in the asymptotic expansion of $s(x)$. With the same model, Alexey et al. [12] find the Mellin transform of the first hitting time of the origin and give an expression for its density.

Here, we show that the cumulative function of the first hitting time for one Lévy process belongs to $\mathcal{C}(\mathbb{R}_+^* \times \mathbb{R}_+^*)$ and for any $x \in \mathbb{R}_+^*$, to $\mathcal{C}(\mathbb{R}_+)$ and then we derive the first known closed form expression which characterizes the law of (τ_x, K_x, L_x) . The paper is organized as follows: Section 2 presents the model and the aim of the paper, Section 3 studies the regularity of the law of τ_x , Section 4 provides the joint law of (τ_x, K_x, L_x) .

2 Model and Problem to solve

On a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, let X be a Lévy process, right continuous with left limit (RCLL) starting at 0. It is defined as

$$X_t = mt + W_t + \sum_{i=1}^{N_t} Y_i \quad (1)$$

where $m \in \mathbb{R}$, W is a standard Brownian motion, N a Poisson process with constant positive intensity λ and $(Y_i, i \in \mathbb{N}^*)$ is a sequence of independent identically distributed random variables with a distribution function F_Y . We suppose

that the following σ -fields $\sigma(N_t, t \geq 0)$, $\sigma(Y_i, i \in \mathbb{N}^*)$ and $\sigma(W_t, t \geq 0)$ are independent. We are interested in the first hitting time at a level $x > 0$,

$$\tau_x := \inf\{t \geq 0, X_t \geq x\}. \quad (2)$$

We also consider the overshoot and the undershoot respectively defined by

$$K_x := X_{\tau_x} - x, \quad (3)$$

$$L_x := x - X_{\tau_x-}. \quad (4)$$

For $\tilde{X}_t := mt + W_t$ and $\tilde{\tau}_x := \inf\{t \geq 0; \tilde{X}_t \geq x\}$, I. Karatzas and S. E. Shreve in [10] shown that the law of $\tilde{\tau}_x$ is of the form $\tilde{f}(u, x)du + \mathbb{P}(\tilde{\tau}_x = \infty)\delta_\infty(du)$ where

$$\tilde{f}(u, x) = \frac{|x|}{\sqrt{2\pi u^3}} \exp\left[-\frac{1}{2u}(x - mu)^2\right] 1_{]0, +\infty[}(u) \text{ and } \mathbb{P}(\tilde{\tau}_x = \infty) = 1 - e^{mx - |mx|}. \quad (5)$$

L. Coutin and D. Dorobantu [5] prove the existence of the density $f_{\tau_x}(t, x)$ of τ_x and show that:

$$f_{\tau_x}(t, x) = \begin{cases} \lambda \mathbb{E}(1_{\tau_x > t}(1 - F_Y)(x - X_t)) + \mathbb{E}(1_{\tau_x > T_{N_t}} \tilde{f}(t - T_{N_t}, x - X_{T_{N_t}})), & \forall t > 0 \\ \frac{\lambda}{2}(2 - F_Y(x) - F_Y(x_-)) + \frac{\lambda}{4}(F_Y(x) - F_Y(x_-)) & \text{if } t = 0 \end{cases} \quad (6)$$

and $\mathbb{P}(\tau_x = \infty) = 0$ if and only if $m + \lambda \mathbb{E}(Y_1) \geq 0$.

For a more general jump-diffusion process, Roynette et al. [16] show that the Laplace transform of (τ_x, K_x, L_x) is solution of some kind of random integral equation.

The problem addressed in this paper is studying the regularity of the density of τ_x on $]0, +\infty[\times]0, +\infty[$, then at $t = 0$ for a strictly positive level x fixed and compute an expression for the joint distribution of the triplet (τ_x, K_x, L_x) .

3 Regularity of f_{τ_x}

This section deals with the regularity. The first subsection 3.1 treats the continuity on $]0, \infty[\times]0, \infty[$ as well as the last one 3.2 studies the regularity with respect to time at 0.

3.1 Regularity of the density f_{τ_x} on $]0, \infty[\times]0, \infty[$

Here, our goal is to prove Proposition 1 which asserts the regularity of τ_x density law on $]0, +\infty[\times]0, +\infty[$.

Proposition 1. *The application defined on $]0, +\infty[\times]0, +\infty[$ by*

$$(t, x) \longrightarrow f_{\tau_x}(t, x) = \lambda \mathbb{E}(1_{\{\tau_x > t\}}(1 - F_Y)(x - X_t)) + \mathbb{E}(1_{\tau_x > T_{N_t}} \tilde{f}(t - T_{N_t}, x - X_{T_{N_t}}))$$

is almost surely continuous.

Proof. Let be $(t_0, x_0) \in \mathbb{R}_+^* \times \mathbb{R}_+^*$. We denote

$$\Omega' = \left\{ \omega \in \Omega \text{ such that } T_{N_{t_0}(\omega)}(\omega) \neq t_0, \tau_{x_0}(\omega) \neq t_0, x_0 - X_{t_0}(\omega) \notin D_{F_Y} \right\}$$

where D_{F_Y} is the set of the points of discontinuity of the distribution function F_Y .

We assert that $\mathbb{P}(\Omega') = 1$: Indeed, we have

$$1 - \mathbb{P}(\Omega') \leq \mathbb{P}(\tau_x = 0) + \mathbb{P}(X_{t_0} = x_0) + \mathbb{P}(T_{N_{t_0}} = t_0) + \mathbb{P}(x_0 - X_{t_0} \in D_{F_Y}).$$

Since τ_{x_0} , $T_{N_{t_0}}$ and X_{t_0} have a densities with respect to the Lebesgue measure and D_{F_Y} is almost countable, it follows

$$\mathbb{P}(\tau_x = 0) = \mathbb{P}(X_{t_0} = x_0) = \mathbb{P}(T_{N_{t_0}} = t_0) = \mathbb{P}(x_0 - X_{t_0} \in D_{F_Y}) = 0.$$

Note that $\mathbf{1}_{\{\tau_x > T_{N_t}\}} = \mathbf{1}_{\{X_{T_{N_t}}^* < x\}}$ where $X_t^* = \sup_{u \leq t} X_u$.

The random variable $X_{T_{N_{t_0}}}^*$ is reached by the process X either before $T_{N_{t_0}}$ either at $T_{N_{t_0}}$.

Let ω be fixed in Ω' .

• On the event $\{\tau_{x_0} > T_{N_{t_0}}\} = \{X_{T_{N_{t_0}}}^* < x_0\}$, $X_{T_{N_{t_0}(\omega)}(\omega)}^* = X_{T_{N_{t_0}(\omega)}(\omega)}(\omega) \neq x_0$.

Thus, if $X_{T_{N_{t_0}(\omega)}(\omega)}^*$ is reached at $T_{N_{t_0}(\omega)}(\omega)$, either it is less than x_0 or more than x_0 .

If $X_{T_{N_{t_0}(\omega)}(\omega)}^*(\omega) = X_{T_{N_{t_0}(\omega)}(\omega)}(\omega) < x_0$, since $T_{N_{t_0}(\omega)}(\omega) < t_0 < T_{N_{t_0}(\omega)+1}(\omega)$, there exists $\varepsilon_0(\omega) > 0$ and $\delta_0(\omega) > 0$ such that for any t satisfying $|t - t_0| \leq \delta_0(\omega)$ we have $x_0 - \varepsilon_0(\omega) < X_{T_{N_t}(\omega)}^*(\omega) < x_0 + \varepsilon_0(\omega)$.

That means for (t, x) such that $|t - t_0| < \delta_0(\omega)$ and $|x - x_0| < \varepsilon_0(\omega)$, we have

$$\mathbf{1}_{\{\tau_x(\omega) > T_{N_t(\omega)}(\omega)\}} = \mathbf{1}_{\{\tau_{x_0}(\omega) > T_{N_{t_0}(\omega)}(\omega)\}} = 1.$$

If $X_{T_{N_{t_0}(\omega)}(\omega)}^*(\omega) = X_{T_{N_{t_0}(\omega)}(\omega)}(\omega) > x_0$, since $T_{N_{t_0}(\omega)}(\omega) < t_0 < T_{N_{t_0}(\omega)+1}(\omega)$, there exists $\varepsilon_1(\omega) > 0$ and $\delta_1(\omega) > 0$ such that for any t satisfying $|t - t_0| \leq \delta_1(\omega)$ we have $x_0 - \varepsilon_1(\omega) < X_{T_{N_t}(\omega)}^* < x_0 + \varepsilon_1(\omega)$.

That means for $|t - t_0| < \delta_1(\omega)$ and $|x - x_0| < \varepsilon_1(\omega)$,

$$\mathbf{1}_{\{\tau_x(\omega) > T_{N_t(\omega)}(\omega)\}} = \mathbf{1}_{\{\tau_{x_0}(\omega) > T_{N_{t_0}(\omega)}(\omega)\}} = 0.$$

In the two above cases, we conclude that for any $\omega \in \Omega$, there exists $\delta(\omega) > 0$ and $\varepsilon(\omega) > 0$ such that:

$$|t - t_0| < \delta(\omega) \text{ and } |x - x_0| < \varepsilon(\omega) \text{ imply that } \mathbf{1}_{\{\tau_x(\omega) > T_{N_t(\omega)}(\omega)\}} = \mathbf{1}_{\{\tau_{x_0}(\omega) > T_{N_{t_0}(\omega)}(\omega)\}} \quad (7)$$

• If $X_{T_{N_{t_0}(\omega)}(\omega)}^*(\omega)$ is reached at $v < T_{N_{t_0}(\omega)}(\omega)$, either it is less than x_0 or more than x_0 .

If $X_{T_{N_{t_0}(\omega)}(\omega)}^*(\omega) = X_v(\omega) < x_0$, since $T_{N_{t_0}(\omega)}(\omega) < t_0 < T_{N_{t_0}(\omega)+1}(\omega)$, there exists $\varepsilon_4(\omega) > 0$ and $\delta_4(\omega) > 0$ such that for any t satisfying $|t - t_0| \leq \delta_4(\omega)$ we have $x_0 - \varepsilon_4(\omega) < X_{T_{N_t}(\omega)}^*(\omega) < x_0 + \varepsilon_4(\omega)$.

That means for (t, x) such that $|t - t_0| < \delta_4(\omega)$ and $|x - x_0| < \varepsilon_4(\omega)$, we have

$$\mathbf{1}_{\{\tau_x > T_{N_t}\}} = \mathbf{1}_{\{\tau_{x_0} > T_{N_{t_0}}\}} = 1.$$

If $X_{T_{N_{t_0}(\omega)}(\omega)}^*(\omega) = X_v(\omega) > x_0$, since $T_{N_{t_0}(\omega)}(\omega) < t_0 < T_{N_{t_0}(\omega)+1}(\omega)$, there exists $\varepsilon_5(\omega) > 0$ and $\delta_5(\omega) > 0$ such that for any t satisfying $|t - t_0| \leq \delta_5(\omega)$ we have $x_0 - \varepsilon_5(\omega) < X_{T_{N_t}}^* < x_0 + \varepsilon_5(\omega)$. That means for $|t - t_0| < \delta_5(\omega)$ and $|x - x_0| < \varepsilon_5(\omega)$,

$$\mathbf{1}_{\{\tau_x > T_{N_t}\}} = \mathbf{1}_{\{\tau_{x_0} > T_{N_{t_0}}\}} = 0.$$

If $X_{T_{N_{t_0}(\omega)}(\omega)}^*(\omega) = X_v(\omega) = x_0$, we consider a function which is equal to $\tilde{f}$ on $]0, +\infty[\times]0, +\infty[$ and 0 on $]0, +\infty[\times \mathbb{R}_-$. We denote it $\tilde{f}$ again. This function is everywhere continuous. Since $T_{N_{t_0}(\omega)}(\omega) < t_0 < T_{N_{t_0}(\omega)+1}(\omega)$, and $(t, x) \longrightarrow \tilde{f}(t - T_{N_t}(\omega), x - X_{T_{N_t}(\omega)}(\omega))$ is continuous at (t_0, x_0) , there exists $\varepsilon_6(\omega) > 0$ and $\delta_6(\omega) > 0$ such that for any (t, x) satisfying $|t - t_0| \leq \delta_6(\omega)$ and $|x - x_0| \leq \varepsilon_6(\omega)$, we have

$$\begin{aligned} & \lim_{(t,x) \rightarrow (t_0,x_0)} \mathbf{1}_{\{X_{T_{N_t}(\omega)}^*(\omega) < x\}} \tilde{f}(t - T_{N_t}(\omega), x - X_{T_{N_t}(\omega)}(\omega)) \quad (8) \\ &= \mathbf{1}_{\{X_{T_{N_{t_0}(\omega)}(\omega)}^* < x_0\}} \tilde{f}(t_0 - T_{N_{t_0}(\omega)}, x_0 - X_{T_{N_{t_0}(\omega)}}(\omega)) = 0. \end{aligned}$$

Using uniform integrability of the family $\left(\mathbf{1}_{\tau_x > T_{N_t}} \tilde{f}(t - T_{N_t}, x - X_{T_{N_t}}), \quad t > 0, \quad x > 0 \right)$, obtained from Lemma 2, the continuity of

$$(t, x) \longrightarrow \mathbb{E} \left(\mathbf{1}_{\tau_x > T_{N_t}} \tilde{f}(t - T_{N_t}, x - X_{T_{N_t}}) \right)$$

at (t_0, x_0) follows.

Since the family $\left((\mathbf{1}_{\{\tau_x > t\}}(1 - F_Y)(x - X_t), t > 0, x > 0) \right)$ is bounded by 1, it is uniformly integrable and we proceed analogously to the previous to obtain the continuity at (t_0, x_0) of

$$(t, x) \longrightarrow \mathbb{E} \left(\mathbf{1}_{\{\tau_x > t\}}(1 - F_Y)(x - X_t) \right).$$

□

We now study the regularity with respect to time at 0.

3.2 Regularity of f_{τ_x} with respect to time at 0

The next two propositions show that for any fixed $x > 0$, the f_{τ_x} density law is continuous with respect to time at 0.

Proposition 2. *Let be $x > 0$ fixed, we have*

$$\lim_{t \rightarrow 0} \mathbb{E}(\mathbf{1}_{\tau_x > t} [1 - F_Y](x - X_t)) = \frac{1}{2} (2 - F_Y(x) - F_Y(x_-)).$$

Proof. We have

$$\mathbb{E}(\mathbf{1}_{\tau_x > t} [1 - F_Y](x - X_t)) = \mathbb{E}(\mathbf{1}_{\{N_t=0\}} \mathbf{1}_{\tau_x > t} [1 - F_Y](x - X_t)) + \mathbb{E}(\mathbf{1}_{\{N_t>0\}} \mathbf{1}_{\tau_x > t} [1 - F_Y](x - X_t)).$$

But

$$\begin{aligned} (i) \quad & 0 \leq \lim_{t \rightarrow 0} \mathbb{E}(\mathbf{1}_{\{N_t>0\}} \mathbf{1}_{\tau_x > t} [1 - F_Y](x - X_t)) \leq \lim_{t \rightarrow 0} \mathbb{P}(N_t \geq 1) = \lim_{t \rightarrow 0} 1 - e^{-at} = 0 \\ (ii) \quad & \mathbb{E}(\mathbf{1}_{\{N_t=0\}} \mathbf{1}_{\tau_x > t} [1 - F_Y](x - X_t)) = e^{-at} \mathbb{E}(\mathbf{1}_{\{\tilde{X}_t^* < x\}} [1 - F_Y](x - \tilde{X}_t)) \end{aligned}$$

We first remark that

$$|\mathbb{E}(\mathbf{1}_{\{\tilde{X}_t^* < x\}} [1 - F_Y](x - \tilde{X}_t)) - \mathbb{E}(\mathbf{1}_{\{\tilde{X}_t < x\}} [1 - F_Y](x - \tilde{X}_t))| \leq |\mathbb{E}(\mathbf{1}_{\{\tilde{X}_t^* < x\}} - \mathbf{1}_{\{\tilde{X}_t < x\}})| = \mathbb{P}(\{\tilde{X}_t < x \leq \tilde{X}_t^*\}).$$

The density function of $(\tilde{X}_t^*, \tilde{X}_t)$ given by Corollary 3.2.1.2 p. 147 [9] yields

$$\mathbb{P}(\{\tilde{X}_t < x \leq \tilde{X}_t^*\}) = \int_x^\infty db \int_{-\infty}^b da \frac{2(2b-a)}{\sqrt{2\pi t^3}} \exp \left[-\frac{(2b-a)^2}{2t} + ma - \frac{m^2}{2}t \right].$$

This integral is bounded (with respect to a multiplicative constant C) by

$$\mathbb{P}(\{\tilde{X}_t < x \leq \tilde{X}_t^*\}) \leq C \int_x^\infty db \int_{-\infty}^b da \frac{(2b-a)}{\sqrt{t^3}} \exp \left[-\frac{(2b-a)^2}{2t} \right].$$

Notice that the application $t \rightarrow \frac{(2b-a)}{\sqrt{t^3}} \exp \left[-\frac{(2b-a)^2}{2t} \right]$ is decreasing to 0 when $t \downarrow 0$. So Lebesgue's monotonous convergence theorem proves that

$$\lim_{t \rightarrow 0} \mathbb{P}(\{\tilde{X}_t < x \leq \tilde{X}_t^*\}) = 0. \quad (9)$$

Secondly,

$$\mathbb{E}(\mathbf{1}_{\{\tilde{X}_t < x\}} [1 - F_Y](x - \tilde{X}_t)) = \mathbb{E}(\mathbf{1}_{\{\tilde{X}_t \leq 0\}} [1 - F_Y](x - \tilde{X}_t)) + \mathbb{E}(\mathbf{1}_{\{0 \leq \tilde{X}_t < x\}} [1 - F_Y](x - \tilde{X}_t)).$$

Since F_Y is bounded and RCLL and $\tilde{X}$ continuous, Lebesgue's dominated convergence theorem yields

$$\lim_{t \rightarrow 0} \mathbb{E}(\mathbf{1}_{\{N_t=0\}} \mathbf{1}_{\tau_x > t} [1 - F_Y](x - X_t)) = \frac{1}{2} (2 - F_Y(x) - F_Y(x_-)).$$

□

Proposition 3. *Let be $x > 0$ fixed, we have*

$$\lim_{t \rightarrow 0} \mathbb{E} \left(\mathbf{1}_{\{\tau_x > T_{N_t}\}} \tilde{f}(t - T_{N_t}, x - X_{T_{N_t}}) \right) = \frac{\lambda}{4} (F_Y(x) - F_Y(x_-)).$$

Proof. (i) We first deal with

$$\lim_{t \rightarrow 0} \mathbb{E} \left(\mathbf{1}_{\{N_t=0, \tau_x > T_{N_t}\}} \tilde{f}(t - T_{N_t}, x - X_{T_{N_t}}) \right) = \lim_{t \rightarrow 0} \mathbb{E}(\mathbf{1}_{\{N_t=0\}} \tilde{f}(t, x)) = \lim_{t \rightarrow 0} (1 - e^{-at}) \tilde{f}(t, x) = 0$$

using Definition (5).

(ii) Then we deal with

$$\mathbb{E} \left(\mathbf{1}_{\{N_t \geq 2, \tau_x > T_{N_t}\}} \tilde{f}(t - T_{N_t}, x - X_{T_{N_t}}) \right).$$

We use Lemma A.1 of [5] for $p = 1$, law of G being the Gaussian law $\mathcal{N}(0, 1)$, then

$$\mathbb{E} \left(\tilde{f}(u, \mu + \sigma G) \mathbf{1}_{\{\mu + \sigma G > 0\}} \right) = \frac{1}{\sqrt{2\pi}} \frac{e^{-\frac{(\mu - mu)^2}{2(\sigma^2 + u)}}}{\sqrt{u}(\sigma^2 + u)} \mathbb{E} \left[\left(\sigma G + \sqrt{\frac{u}{\sigma^2 + u}} (\mu - mu) + m \sqrt{u(\sigma^2 + u)} \right)_+ \right].$$

Using $C_{1/2} = \sup \sqrt{y} e^{-\frac{y^2}{2}}$, $y = \frac{\mu - mu}{\sqrt{\sigma^2 + u}}$,

$$\mathbb{E} \left(\tilde{f}(u, \mu + \sigma G) \mathbf{1}_{\{\mu + \sigma G > 0\}} \right) \leq \frac{1}{\sqrt{2\pi}} \left[\frac{1}{\sqrt{u}(\sigma^2 + u)} \left(\mathbb{E}|\sigma G| + \sqrt{u} C_{1/2} + |m| \sqrt{u(\sigma^2 + u)} \right) \right].$$

For $u = t - T_{N_t}$, $\sigma^2 = T_{N_t}$ and $u + \sigma^2 = t$, we obtain using the independence between the Poisson process and the Brownian motion

$$\mathbb{E} \left(\mathbf{1}_{\{N_t \geq 2\}} \mathbf{1}_{\{\tau_x > T_{N_t}\}} \tilde{f}(t - T_{N_t}, x - X_{T_{N_t}}) \right) \leq \mathbb{E} \left(\mathbf{1}_{\{N_t \geq 2\}} \frac{1}{t \sqrt{2\pi}} \left[\frac{E[|\sigma G|]}{\sqrt{t - T_{N_t}}} + C_{1/2} + |m| \sqrt{t} \right] \right).$$

So, using $\mathbb{P}(\{N_t \geq 2\}) = 0(t^2)$ and Lemma 1,

$$\lim_{t \rightarrow 0} \mathbb{E} \left(\mathbf{1}_{\{N_t \geq 2\}} \mathbf{1}_{\{\tau_x > T_{N_t}\}} \tilde{f}(t - T_{N_t}, x - X_{T_{N_t}}) \right) = 0.$$

(iii) Finally we deal with

$$A_t = \mathbb{E} \left(\mathbf{1}_{\{N_t=1, \tau_x > T_{N_t}\}} \tilde{f}(t - T_{N_t}, x - X_{T_{N_t}}) \right) = \mathbb{E} \left(\mathbf{1}_{\{N_t=1, \tau_x > T_1\}} \tilde{f}(t - T_1, x - X_{T_1}) \right).$$

Since the event $\{N_t = 1, \tau_x > T_1\} = \{T_1 \leq t < T_2, \tilde{X}_{T_1}^* < x, \tilde{X}_{T_1} + Y_1 < x\}$ we have

$$A_t = \mathbb{E} \left(\mathbf{1}_{\{T_1 \leq t < T_2, \tilde{X}_{T_1}^* < x, \tilde{X}_{T_1} + Y_1 < x\}} \tilde{f}(t - T_1, x - \tilde{X}_{T_1} - Y_1) \right).$$

Using the law of $T_2 - T_1$ and its independence from $T_1, \tilde{X}_{T_1}^*, \tilde{X}_{T_1}, Y_1$, it follows that

$$A_t = \mathbb{E} \left(e^{-a(t-T_1)} \mathbf{1}_{\{T_1 \leq t, \tilde{X}_{T_1}^* < x, \tilde{X}_{T_1} + Y_1 < x\}} \tilde{f}(t - T_1, x - \tilde{X}_{T_1} - Y_1) \right)$$

Using the law of T_1 and the independence between T_1 and $(\tilde{X}, Y)$, yields

$$A_t = ae^{-at} \int_0^t du \mathbb{E} \left(\mathbf{1}_{\{\tilde{X}_u^* < x, \tilde{X}_u + Y_1 < x\}} \tilde{f}(t-u, x - \tilde{X}_u - Y_1) \right).$$

Since $\tilde{X}_u$ and Y_1 are independent, conditioning by $(\tilde{X}_u, Y_1)$ and using Lemma 4 for $c = x$,

$$A_t = ae^{-at} \int_0^t du \mathbb{E} \left(\mathbf{1}_{\{\tilde{X}_u < \min(x, x - Y_1)\}} \left[1 - e^{-\frac{2x^2 - 2x\tilde{X}_u}{u}} \right] f(t-u, x - \tilde{X}_u - Y_1) \right).$$

The change of variable $u = ts$ leads to

$$A_t = ae^{-at} \int_0^1 tds \mathbb{E} \left(\mathbf{1}_{\{\tilde{X}_{ts} < \min(x, x - Y_1)\}} \left[1 - e^{-\frac{2x^2 - 2x\tilde{X}_{ts}}{ts}} \right] f(t(1-s), x - \tilde{X}_{ts} - Y_1) \right).$$

The density of $\tilde{X}_{ts} : \frac{1}{\sqrt{2\pi ts}} e^{-\frac{(g-mts)^2}{2ts}}$ and $\tilde{f}$ defined in (5) yield

$$A_t = ae^{-at} \int_0^1 tds \mathbb{E} \left(\int_{-\infty}^{\min(x, x - Y_1)} \left[1 - e^{-\frac{2x^2 - 2xg}{ts}} \right] \frac{x - Y_1 - g}{\sqrt{2\pi t^3(1-s)^3}} e^{-\frac{(x-g-Y_1-mt(1-s))^2}{2t(1-s)}} \frac{1}{\sqrt{2\pi ts}} e^{-\frac{(g-mts)^2}{2ts}} dg \right).$$

Let be $z = x - Y_1 - mt(1-s)$, $y = mts$, $u = t(1-s)$ and $v = ts$, with

$$\begin{aligned} \frac{vz + uy}{u + v} &= \frac{ts[x - Y_1 - mt(1-s)] + t(1-s)mts}{t} = s(x - Y_1), \\ z - y &= x - Y_1 - mt(1-s) - mts = x - Y_1 - mt. \end{aligned}$$

By Lemma 3,

$$A_t = ae^{-at} \int_0^1 tds \mathbb{E} \left(\int_{-\infty}^{\min(x, x - Y_1)} \left[1 - e^{-\frac{2x^2 - 2xg}{ts}} \right] \frac{x - Y_1 - g}{\sqrt{(2\pi)^2 t^4 (1-s)^3 s}} e^{-\frac{(g-s(x-Y_1))^2}{2ts(1-s)} - \frac{(x-Y_1-mt)^2}{2t^2 s(1-s)}} dg \right).$$

A new change of variable $g' = \frac{g-s(x-Y_1)}{\sqrt{ts(1-s)}}$ meaning $g = \sqrt{ts(1-s)}g' + s(x - Y_1)$, and $x - Y_1 - g = (x - Y_1)(1-s) - \sqrt{ts(1-s)}g'$ implies

$$A_t = ae^{-at} \int_0^1 ds \int dg' \sqrt{t^3 s(1-s)} \mathbb{E} \left(\mathbf{1}_{\{g' < \frac{\min(x, x - Y_1) - s(x - Y_1)}{\sqrt{ts(1-s)}}\}} \left[1 - e^{-\frac{2x^2 - 2x(\sqrt{ts(1-s)}g' + s(x - Y_1))}{ts}} \right] \frac{(x - Y_1)(1-s) - \sqrt{ts(1-s)}g'}{\sqrt{(2\pi)^2 t^4 (1-s)^3 s}} e^{-\frac{(g')^2}{2} - \frac{(x - Y_1 - mt)^2}{2t^2 s(1-s)}} \right).$$

This would mean

$$A_t = ae^{-at} \int_0^1 ds \int dg' \mathbb{E} \left[\mathbf{1}_{\{g' < \frac{\min(x, x-Y_1)-s(x-Y_1)}{\sqrt{ts(1-s)}}\}} \left[1 - e^{-\frac{2x^2-2x(\sqrt{ts(1-s)}g'+s(x-Y_1))}{ts}} \right] e^{-\frac{(g')^2}{2}-\frac{(x-Y_1-mt)^2}{2t^2s(1-s)}} \left[\frac{(x-Y_1)}{2\pi\sqrt{t}} - \frac{g'\sqrt{s}}{2\pi\sqrt{1-s}} \right] \right].$$

On the set $\{g' < c\}$, $2x^2 - 2xc > 0$. This would mean $\{g' < \frac{\min(x, x-Y_1)-s(x-Y_1)}{\sqrt{ts(1-s)}}\}$, where

$$c = \frac{x-Y_1-s(x-Y_1)}{\sqrt{ts(1-s)}} - \frac{2x^2-2x(\sqrt{ts(1-s)}g'+s(x-Y_1))}{ts} < 0, \text{ and}$$

$$\left| 1 - e^{-\frac{2x^2-2x(\sqrt{ts(1-s)}g'+s(x-Y_1))}{ts}} \right| \mathbf{1}_{\{g' < \frac{\min(x, x-Y_1)-s(x-Y_1)}{\sqrt{ts(1-s)}}\}} \leq 1;$$

with

$$\lim_{t \rightarrow 0} \left[1 - e^{-\frac{2x^2-2x(\sqrt{ts(1-s)}g'+s(x-Y_1))}{ts}} \right] \mathbf{1}_{\{g' < \frac{\min(x, x-Y_1)-s(x-Y_1)}{\sqrt{ts(1-s)}}\}} = 1.$$

Moreover,

$$\begin{aligned} \frac{|x-Y_1|}{\sqrt{t}} e^{-\frac{(x-Y_1-mt)^2}{2t^2s(1-s)}} &\leq \frac{|mt| + |x-Y_1-mt|}{\sqrt{t}} e^{-\frac{(x-Y_1-mt)^2}{2t^2s(1-s)}} \\ &\leq \frac{|mt| + C_{1/2}\sqrt{t^2s(1-s)}}{\sqrt{t}} = \sqrt{t} (|m| + C_{1/2}\sqrt{s(1-s)}) \end{aligned}$$

Thus

$$\lim_{t \rightarrow 0} \frac{|x-Y_1|}{\sqrt{t}} e^{-\frac{(x-Y_1-mt)^2}{2t^2s(1-s)}} = 0.$$

Finally,

$$e^{-\frac{(g')^2}{2}-\frac{(x-Y_1-mt)^2}{2t^2s(1-s)}} \frac{|g'|\sqrt{s}}{2\pi\sqrt{1-s}} \leq e^{-\frac{(g')^2}{2}} \frac{|g'|\sqrt{s}}{2\pi\sqrt{1-s}}$$

and

$$\lim_{t \rightarrow 0} \mathbf{1}_{\{g' < \frac{\min(x, x-Y_1)-s(x-Y_1)}{\sqrt{ts(1-s)}}\}} e^{-\frac{(g')^2}{2}-\frac{(x-Y_1-mt)^2}{2t^2s(1-s)}} \frac{g'\sqrt{s}}{2\pi\sqrt{1-s}} = e^{-\frac{(g')^2}{2}} \frac{g'\sqrt{s}}{2\pi\sqrt{1-s}} \mathbf{1}_{\{Y_1=0\}} \mathbf{1}_{\{g'<0\}}.$$

By Lebesgue's dominated convergence theorem ,

$$\lim_{t \rightarrow 0} A_t = -a\mathbb{P}(Y_1 = x) \int_0^1 ds \int_{\{g'<0\}} dg' e^{-\frac{(g')^2}{2}} \frac{g'\sqrt{s}}{2\pi\sqrt{1-s}}$$

and $\lim_{t \rightarrow 0} A_t = a\mathbb{P}(Y_1 = x) \int_0^1 \frac{\sqrt{s}}{2\pi\sqrt{1-s}} ds = \frac{a\mathbb{P}(Y=x)}{4}$. Indeed, $\beta(3/2, 1/2) = \frac{1}{2}\gamma(1/2)^2$ and $\gamma(1/2) = \pi$. $\square$

4 The joint law

If the default time coincides with a jump time of the process X , it is also important to have information on the deficit, namely overshoot, right after the default and on the surplus, namely undershoot of the firm, immediately before the default time. Therefore, A. Volpi et al. [16] deal with the asymptotic behavior of the triplet (τ_x, K_x, L_x) by showing after a renormalization of τ_x that it converges in distribution as x goes to ∞ . To characterize the joint law of (τ_x, K_x, L_x) on $\mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+$ for any $x > 0$, this section's main result is the next theorem. From now on, we consider two continuous bounded functions Φ and Ψ . The methodology used for the proof is inspired from Coutin-Dorobantu [5] who study the law of τ_x which is here the first marginal distribution. It consists in splitting $\mathbb{E}(1_{t < \tau_x \leq t+h} \Phi(K_x) \Psi(L_x))$ following the values of $N_{t+h} - N_t$ for $t = 0$ then for $t > 0$.

Theorem 1. *The joint law of the triplet (τ_x, K_x, L_x) , conditionally on $\{\tau_x < \infty\}$, is given on $\mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+$ by $p(\cdot, \cdot, \cdot, x)$ such that:*

$$\begin{aligned} p(0, dk, dl) &= \frac{\lambda}{4} [F_Y(x) - F_Y(x_-)] \delta_{\{0,0,0\}}(dt, dk, dl) + \lambda F_l(dk) \delta_{\{0,x\}}(dt, dl) \\ &\quad + \frac{\lambda}{2} \Delta F_Y(x) \delta_{\{0,0,x\}}(dt, dk, dl) \end{aligned}$$

and for every $t > 0$,

$$\begin{aligned} p(dt, dk, dl) &= \mathbb{E}[\mathbf{1}_{\{\tau_x > T_{N_t}\}} \tilde{f}(t - T_{N_t}, x - X_{T_{N_t}})] \delta_{\{0,0\}}(dk, dl) dt \\ &\quad + \lambda \mathbb{E} \left[\mathbf{1}_{\{k \geq 0, l \geq 0\}} \mathbf{1}_{\{\tau_x > T_{N_t}\}} f_0(x - X_{T_{N_t}} - l) \right] F_l(dk) dl dt \\ &\quad - \lambda \mathbb{E} \left[\mathbf{1}_{\{k \geq 0, l \geq 0\}} \mathbf{1}_{\{\tau_x > T_{N_t}\}} f_0(X_{T_{N_t}} - x - l) \exp(2m(x - X_{T_{N_t}})) \right] F_l(dk) dl dt \end{aligned}$$

where f_0 is the density function of a Gaussian random variable with mean $\mu = m(t - T_{N_t})$ and variance $\sigma^2 = t - T_{N_t}$, $\tilde{f}$ is defined by (5), $F_l(dk)$ is the image of $F_Y(dk)$ by the map $y \mapsto y - l$ and $\Delta F_Y(x) = F_Y(x) - F_Y(x_-)$.

Remark 1. Referring to [16], for all $x > 0$, the first passage time τ_x is finite almost surely if and only if $m + \mathbb{E}(Y_1) \geq 0$.

To prove the theorem, we use Propositions 4 and 5. Indeed, Proposition 4 gives the law at time $t = 0$ and Proposition 5 deals with time $t > 0$. The proof of Proposition 5 is broken in three parts: Step 1, Step 2 and Proposition 6.

Proposition 4.

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{1}{h} \mathbb{E}(1_{\tau_x \leq h} \Phi(K_x) \Psi(L_x)) &= \Phi(0) \Psi(0) \frac{\lambda}{4} [F_Y(x) - F_Y(x_-)] + \lambda \mathbb{E}[\Phi(Y_1 - x) \Psi(x) \mathbf{1}_{\{Y_1 > x\}}] \\ &\quad + \frac{\lambda}{2} \mathbb{E}[\Phi(0) \Psi(Y_1) \mathbf{1}_{\{Y_1 = x\}}]. \end{aligned}$$

Proof. We split $\mathbb{E}(1_{\tau_x \leq h} \Phi(K_x) \Psi(L_x))$ according to the values of N_h :

$$\begin{aligned} \mathbb{E}(1_{\tau_x \leq h} \Phi(K_x) \Psi(L_x)) &= \mathbb{E}(1_{\tau_x \leq h} \mathbf{1}_{\{N_h=0\}} \Phi(K_x) \Psi(L_x)) + \mathbb{E}(1_{\tau_x \leq h} \mathbf{1}_{\{N_h=1\}} \Phi(K_x) \Psi(L_x)) \\ &\quad + \mathbb{E}(1_{\tau_x \leq h} \mathbf{1}_{\{N_h \geq 2\}} \Phi(K_x) \Psi(L_x)). \end{aligned}$$

By hypothesis, Φ and Ψ are bounded and on the event $\{N_h = 0\}$, the law of τ_x is the one of $\tilde{\tau}_x$, so has the continuous density $\tilde{f}(\cdot, x)$ defined in (5). Since

$$\mathbb{E}(1_{\tau_x \leq h} \mathbf{1}_{\{N_h=0\}} \Phi(K_x) \Psi(L_x)) = \Phi(0) \Psi(0) \mathbb{P}(\tilde{\tau}_x \leq h)$$

and

$$|\mathbb{E}(1_{\tau_x \leq h} \mathbf{1}_{\{N_h \geq 2\}} \Phi(K_x) \Psi(L_x))| \leq \|\Phi\|_\infty \|\Psi\|_\infty \mathbb{P}(N_h \geq 2),$$

it follows that

$$\lim_{h \rightarrow 0} \frac{1}{h} \mathbb{E}(1_{\tau_x \leq h} \mathbf{1}_{\{N_h=0\}} \Phi(K_x) \Psi(L_x)) = 0 \quad (10)$$

and

$$\lim_{h \rightarrow 0} \frac{1}{h} \mathbb{E}(1_{\tau_x \leq h} \mathbf{1}_{\{N_h \geq 2\}} \Phi(K_x) \Psi(L_x)) = 0.$$

It remains to study $\mathbb{E}(1_{\tau_x \leq h} \mathbf{1}_{\{N_h=1\}} \Phi(K_x) \Psi(L_x))$. For this purpose, we split it as in Coutin and Dorobantu [5] according to the relative positions of τ_x and T_1 the first jump time of the process N .

$$\begin{aligned} \mathbb{E}(1_{\tau_x \leq h} \mathbf{1}_{\{N_h=1\}} \Phi(K_x) \Psi(L_x)) &= \mathbb{E}(1_{\tau_x \leq h} \mathbf{1}_{\{N_h=1\}} \mathbf{1}_{\{\tau_x < T_1\}} \Phi(K_x) \Psi(L_x)) \\ &\quad + \mathbb{E}(1_{\tau_x \leq h} \mathbf{1}_{\{N_h=1\}} \mathbf{1}_{\{\tau_x = T_1\}} \Phi(K_x) \Psi(L_x)) \\ &\quad + \mathbb{E}(1_{\tau_x \leq h} \mathbf{1}_{\{N_h=1\}} \mathbf{1}_{\{\tau_x > T_1\}} \Phi(K_x) \Psi(L_x)) \\ &= A_1(h) + A_2(h) + A_3(h). \end{aligned}$$

On the set $\{\tau_x \leq h, \tau_x \neq T_1, N_h = 1\}$, the process X is continuous at τ_x . and $K_x = L_x = 0$. Therefore, Step 1 and Step 3 of Subsection 2.1 in [5] imply that:

$$\lim_{h \rightarrow 0} \frac{1}{h} A_1(h) = 0$$

and

$$\lim_{h \rightarrow 0} \frac{1}{h} \mathbb{E}(1_{\tau_x \leq h} \mathbf{1}_{\{N_h=1\}} \mathbf{1}_{\{\tau_x > T_1\}}) = \frac{\lambda}{4} [F_Y(x) - F_Y(x_-)].$$

To study $A_2(h)$, we observe that:

$$\begin{aligned} \tau_x = T_1 \text{ if and only if } \tilde{X}_{T_1}^* < x \text{ and } \tilde{X}_{T_1} + Y_1 > x \\ \text{and on this set } K_x = \tilde{X}_{T_1} + Y_1 - x, \quad L_x = x - \tilde{X}_{T_1} \text{ and } Y_1 > 0. \end{aligned}$$

Therefore ,

$$A_2(h) = \mathbb{E} \left(\mathbf{1}_{\{T_1 \leq h < T_2\}} \mathbf{1}_{\{\tilde{X}_{T_1}^* < x, \tilde{X}_{T_1} + Y > x\}} \Phi(\tilde{X}_{T_1} + Y - x) \Psi(x - \tilde{X}_{T_1}) \right).$$

The independence of $(S_i, i \geq 1)$ and $(Y_1, \tilde{X}, \tilde{\tau}_x)$ leads after integrating with respect to S_2 , then S_1 to

$$\frac{1}{h}A_2(h) = \frac{\lambda e^{-\lambda h}}{h} \int_0^h \mathbb{E} \left(\mathbf{1}_{\{\tilde{X}_u^* < x < \tilde{X}_u + Y_1\}} \Phi(\tilde{X}_u + Y_1 - x) \Psi(x - \tilde{X}_u) \right) du$$

$$|\mathbb{E} \left(\mathbf{1}_{\{\tilde{X}_u^* < x < \tilde{X}_u + Y_1\}} \Phi(\tilde{X}_u + Y_1 - x) \Psi(x - \tilde{X}_u) \right) - \mathbb{E} \left(\mathbf{1}_{\{\tilde{X}_u < x < \tilde{X}_u + Y_1\}} \Phi(\tilde{X}_u + Y_1 - x) \Psi(x - \tilde{X}_u) \right)|$$

is less than $\|\Phi\|_\infty \|\Psi\|_\infty \mathbb{P}(\tilde{X}_u < x < \tilde{X}_u^*)$ which, by (9), goes to zero when u goes to zero. Hence, we obtain

$$\lim_{h \rightarrow 0} \frac{1}{h} A_2(h) = \lim_{h \rightarrow 0} \frac{\lambda e^{-\lambda h}}{h} \int_0^h \mathbb{E} \left(\mathbf{1}_{\{\tilde{X}_u < x < \tilde{X}_u + Y_1\}} \Phi(\tilde{X}_u + Y_1 - x) \Psi(x - \tilde{X}_u) \right) du.$$

But, we have the equality of the sets

$$\{\tilde{X}_u < x < \tilde{X}_u + Y_1\} = \left\{ \{\tilde{X}_u < x < \tilde{X}_u + Y_1\} \cap \{Y_1 \neq x\} \right\} \cup \left\{ \{\tilde{X}_u < x < \tilde{X}_u + Y_1\} \cap \{Y_1 = x\} \right\}.$$

It follows that

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{1}{h} A_2(h) &= \lim_{h \rightarrow 0} \frac{\lambda e^{-\lambda h}}{h} \int_0^h \mathbb{E} \left(\mathbf{1}_{\{\{\tilde{X}_u < x < \tilde{X}_u + Y_1\} \cap \{Y_1 \neq x\}\}} \Phi(\tilde{X}_u + Y_1 - x) \Psi(x - \tilde{X}_u) \right) du \\ &\quad + \lim_{h \rightarrow 0} \frac{\lambda e^{-\lambda h}}{h} \int_0^h \mathbb{E} \left(\mathbf{1}_{\{\{\tilde{X}_u < x < \tilde{X}_u + Y_1\} \cap \{Y_1 = x\}\}} \Phi(\tilde{X}_u + Y_1 - x) \Psi(x - \tilde{X}_u) \right) du. \end{aligned}$$

Since Φ and Ψ are Borel and continuous bounded functions and $\tilde{X}$ continuous, Lebesgue's dominated convergence theorem yields

$$\lim_{h \rightarrow 0} \frac{1}{h} A_2(h) = \lambda \mathbb{E}[\Phi(Y_1 - x) \Psi(x) \mathbf{1}_{\{Y_1 > x\}}] + \frac{\lambda}{2} \mathbb{E}[\Phi(0) \Psi(Y_1) \mathbf{1}_{\{Y_1 = x\}}].$$

□

Proposition 5. *The joint law of the triplet (τ_x, K_x, L_x) , conditionally on $\{\tau_x < \infty\}$, is defined on $\mathbb{R}_+^* \times \mathbb{R}_+ \times \mathbb{R}_+$ as following*

$$\begin{aligned} p(dt, dk, dl) &= \mathbb{E}(1_{\tau_x > T_{N_t}} \tilde{f}(t - T_{N_t}, x - X_{T_{N_t}})) \delta_{\{0,0\}}(dk, dl) dt \\ &\quad + \mathbf{1}_{\{k \geq 0, l \geq 0\}} \mathbb{E} \left[\mathbf{1}_{\{\tau_x > T_{N_t}\}} \frac{\lambda e^{-\frac{(x-l-X_{T_{N_t}}-m(t-T_{N_t}))^2}{2(t-T_{N_t})}}}{\sqrt{2\pi(t-T_{N_t})}} \right] F_l(dk) dl dt \\ &\quad - \mathbf{1}_{\{k \geq 0, l \geq 0\}} \mathbb{E} \left[\mathbf{1}_{\{\tau_x > T_{N_t}\}} \frac{\lambda e^{-\frac{(x-X_{T_{N_t}}+l+m(t-T_{N_t}))^2}{2(t-T_{N_t})} + 2m(x-X_{T_{N_t}})}}}{\sqrt{2\pi(t-T_{N_t})}} \right] F_l(dk) dl dt \end{aligned}$$

where $F_l(dk)$ is the image of $F_Y(dk)$ by the map $y \mapsto y - l$.

Proof. We calculate

$$\lim_{h \rightarrow 0} \frac{1}{h} \mathbb{E}(1_{t < \tau_x \leq t+h} \Phi(K_x) \Psi(L_x)) \quad (11)$$

for $t > 0$ fixed. For this purpose, we split $\mathbb{E}(1_{t < \tau_x \leq t+h} \Phi(K_x) \Psi(L_x))$ according to the values of $N_{t+h} - N_t$ as following:

$$\begin{aligned} \mathbb{E}(1_{t < \tau_x \leq t+h} \Phi(K_x) \Psi(L_x)) &= \mathbb{E}(1_{t < \tau_x \leq t+h} \Phi(K_x) \Psi(L_x) 1_{N_{t+h} - N_t = 0}) \\ &\quad + \mathbb{E}(1_{t < \tau_x \leq t+h} \Phi(K_x) \Psi(L_x) 1_{N_{t+h} - N_t = 1}) \\ &\quad + \mathbb{E}(1_{t < \tau_x \leq t+h} \Phi(K_x) \Psi(L_x) 1_{N_{t+h} - N_t \geq 2}). \end{aligned} \quad (12)$$

So we deal the proof with three steps, the third one being Proposition 6

(i) The third term of the right hand side of (12) is upper bounded as following:

$$\mathbb{E}(1_{t < \tau_x \leq t+h} \Phi(K_x) \Psi(L_x) 1_{N_{t+h} - N_t \geq 2}) \leq (1 - e^{ah} - ahe^{ah}) \|\Phi\|_\infty \|\Psi\|_\infty.$$

Therefore

$$\lim_{h \rightarrow 0} \frac{1}{h} \mathbb{E}(1_{t < \tau_x \leq t+h} \Phi(K_x) \Psi(L_x) 1_{N_{t+h} - N_t \geq 2}) = 0.$$

(ii) Let us study the first term on the right hand side of (12). On the set

$$\{\omega, \quad N_{t+h}(\omega) - N_t(\omega) = 0\}, \text{ we have } L_x = 0 = K_x.$$

Then

$$\mathbb{E}(1_{t < \tau_x \leq t+h} \Phi(K_x) \Psi(L_x) 1_{N_{t+h} - N_t = 0}) = \Phi(0) \Psi(0) \mathbb{E}(1_{t < \tau_x \leq t+h} 1_{N_{t+h} - N_t = 0}).$$

Refer to Equation (4) et seq at Section (2.2) in [5], we have

$$\lim_{h \rightarrow 0} \frac{1}{h} \mathbb{E}(1_{t < \tau_x \leq t+h} 1_{N_{t+h} - N_t = 0}) = \mathbb{E}(1_{\tau_x > T_{N_t}} f(t - T_{N_t}, x - X_{T_{N_t}})).$$

So, we deduce that

$$\lim_{h \rightarrow 0} \frac{1}{h} \mathbb{E}(1_{t < \tau_x \leq t+h} \Phi(K_x) \Psi(L_x) 1_{N_{t+h} - N_t = 0}) = \Phi(0) \Psi(0) \mathbb{E}(1_{\tau_x > T_{N_t}} f(t - T_{N_t}, x - X_{T_{N_t}})).$$

The third step is Proposition 6 which deals with the middle term in (12).

Proposition 6. When h goes to 0,

$$h \mapsto \frac{1}{h} \mathbb{E} [1_{t < \tau_x \leq t+h} \Phi(K_x) \Psi(L_x) 1_{N_{t+h} - N_t = 1}]$$

converges to

$$\begin{aligned} &\int_0^\infty \int_0^{+\infty} \Phi(k) \Psi(l) \mathbb{E} \left[\mathbf{1}_{\{\tau_x > T_{N_t}\}} \frac{\lambda e^{(-\frac{(x-l-X_{T_{N_t}}-m(t-T_{N_t}))^2}{2(t-T_{N_t}})})}{\sqrt{2\pi(t-T_{N_t})}} \right] F_l(dk) dl \\ &- \int_0^{+\infty} \int_0^{+\infty} \Phi(k) \Psi(l) \mathbb{E} \left[\mathbf{1}_{\{\tau_x > T_{N_t}\}} \frac{\lambda e^{(-\frac{(x-X_{T_{N_t}}+l+m(t-T_{N_t}))^2}{2(t-T_{N_t}})+2m(x-X_{T_{N_t}})})}{\sqrt{2\pi(t-T_{N_t})}} \right] F_l(dk) dl \end{aligned}$$

where $F_l(dk)$ is the image of $F_Y(dk)$ by the map $y \mapsto y - l$.

Proof. We split the middle term of (12) according to the values of N_t . Since $\{N_t = n\} = \{T_n \leq t < T_{n+1}\}$ so

$$\begin{aligned} \mathbb{E}(1_{t < \tau_x \leq t+h} \Phi(K_x) \Psi(L_x) 1_{N_{t+h}-N_t=1}) &= \mathbb{E}(1_{t < \tau_x \leq t+h} \Phi(K_x) \Psi(L_x) 1_{T_{N_t} \leq t < T_{N_t+1} \leq t+h < T_{N_t+2}}) \\ &= \sum_{n \geq 0} \mathbb{E}(1_{t < \tau_x \leq t+h} \Phi(K_x) \Psi(L_x) 1_{T_n \leq t < T_{n+1} \leq t+h < T_{n+2}}). \end{aligned} \quad (13)$$

We split again the right term of (13) according to the relative positions between τ_x and T_{n+1} . It follows

$$\begin{aligned} \mathbb{E}(1_{t < \tau_x \leq t+h} \Phi(K_x) \Psi(L_x) 1_{N_{t+h}-N_t=1}) &= \sum_{n \geq 0} \mathbb{E}(\Phi(K_x) \Psi(L_x) 1_{T_n \leq t < \tau_x < T_{n+1} \leq t+h < T_{n+2}}) \\ &\quad + \sum_{n \geq 0} \mathbb{E}(\Phi(K_x) \Psi(L_x) 1_{T_n \leq t < \tau_x = T_{n+1} \leq t+h < T_{n+2}}) \\ &\quad + \sum_{n \geq 0} \mathbb{E}(\Phi(K_x) \Psi(L_x) 1_{T_n \leq t < T_{n+1} < \tau_x \leq t+h < T_{n+2}}). \end{aligned} \quad (14)$$

We recall

$$T_1 = S_1, \quad T_2 = S_1 + S_2, \quad \dots, \quad T_n = \sum_{i=1}^n S_i$$

where $(S_i)_{i \in \mathbb{N}^*}$ is a sequence of independent random variables following an exponential law with parameter λ .

Let be

$$\begin{aligned} A_h^1 &= \sum_{n \geq 0} \mathbb{E}(\Phi(K_x) \Psi(L_x) 1_{T_n \leq t < \tau_x < T_{n+1} \leq t+h < T_{n+2}}) \\ A_h^2 &= \sum_{n \geq 0} \mathbb{E}(\Phi(K_x) \Psi(L_x) 1_{T_n \leq t < \tau_x = T_{n+1} \leq t+h < T_{n+2}}) \\ A_h^3 &= \sum_{n \geq 0} \mathbb{E}(\Phi(K_x) \Psi(L_x) 1_{T_n \leq t < T_{n+1} < \tau_x \leq t+h < T_{n+2}}). \end{aligned}$$

STEP 1: Here, we refer to the analysis of $\frac{1}{h}B_1(h)$, Equation (4) et seq. in [5].

On the sets $\{T_n \leq t < \tau_x < T_{n+1}\}$ and $\{T_{n+1} < \tau_x \leq t+h < T_{n+2}\}$, we have $K_x = 0 = L_x$. So

$$\begin{aligned} A_h^1 &= \Phi(0) \Psi(0) \sum_{n \geq 0} \mathbb{E}(1_{T_n \leq t < \tau_x < T_{n+1} \leq t+h < T_{n+2}}) \\ &= \Phi(0) \Psi(0) \sum_{n \geq 0} \mathbb{E}(1_{T_n \leq t < \tau_x < T_n + S_{n+1} \leq t+h < T_n + S_{n+1} + S_{n+2}}). \end{aligned}$$

Strong Markov property at T_n yields

$$A_h^1 = \Phi(0)\Psi(0) \sum_{n \geq 0} \mathbb{E} \left(\mathbf{1}_{\{T_n \leq t\}} \mathbf{1}_{\{\tau_x > T_n\}} \mathbb{E}^{T_n} (\mathbf{1}_{\{t-T_n < \tilde{\tau}_{x-X_{T_n}} < S_{n+1} < t+h-T_n < S_{n+1}+S_{n+2}\}}) \right)$$

where $\mathbb{E}^{T_n}(\cdot) = \mathbb{E}(\cdot | \mathcal{F}_{T_n})$. Since S_{n+2} is independent from S_{n+1} , $\tau_{x-X_{T_n}}$ and T_n , we obtain

$$\begin{aligned} A_h^1 &= \Phi(0)\Psi(0) \sum_{n \geq 0} \mathbb{E} \left(\mathbf{1}_{\{T_n \leq t\}} \mathbf{1}_{\{\tau_x > T_n\}} \mathbb{E}^{T_n} (e^{-\lambda(t+h-T_n-S_{n+1})} \mathbf{1}_{\{t-T_n < \tilde{\tau}_{x-X_{T_n}} < S_{n+1} < t+h-T_n\}}) \right) \\ &\leq e^{-\lambda h} \Phi(0)\Psi(0) \sum_{n \geq 0} \mathbb{E} \left(\mathbf{1}_{\{\tau_x > T_n\}} \mathbb{E}^{T_n} (e^{\lambda S_{n+1}} \mathbf{1}_{\{t-T_n < \tilde{\tau}_{x-X_{T_n}} < S_{n+1} < t+h-T_n\}}) \right). \end{aligned}$$

The random variables $\tilde{\tau}_{x-X_{T_n}}$ and S_{n+1} are independent and their laws admit a density. Therefore

$$\begin{aligned} \mathbb{E}^{T_n} (e^{\lambda S_{n+1}} \mathbf{1}_{\{t-T_n < \tilde{\tau}_{x-X_{T_n}} < S_{n+1} < t+h-T_n\}}) &= \int_{t-T_n}^{t-T_n+h} \tilde{f}(u, x - X_{T_n}) \int_u^{t-T_n+h} \lambda ds du \\ &= \int_{t-T_n}^{t-T_n+h} \lambda [t - T_n + h - u] \tilde{f}(u, x - X_{T_n}) du. \end{aligned}$$

The change of variable $v = u - (t - T_n)$ implies

$$\mathbb{E}^{T_n} (e^{\lambda S_{n+1}} \mathbf{1}_{\{t-T_n < \tilde{\tau}_{x-X_{T_n}} < S_{n+1} < t+h-T_n\}}) = \int_0^h \lambda [h - v] \tilde{f}(t - T_n + v, x - X_{T_n}) dv.$$

Thus,

$$\begin{aligned} A_h^1 &\leq \lambda e^{-\lambda h} \Phi(0)\Psi(0) \sum_{n \geq 0} \int_0^h \mathbb{E} \left(\mathbf{1}_{\{\tau_x > T_n\}} [h - v] \tilde{f}(t - T_n + v, x - X_{T_n}) \right) dv \\ &\leq \lambda e^{-\lambda h} \Phi(0)\Psi(0) \int_0^h \mathbb{E} \left(\mathbf{1}_{\{\tau_x > T_{N_t}\}} [h - v] \tilde{f}(t - T_{N_t} + v, x - X_{T_{N_t}}) \right) dv. \end{aligned}$$

Similarly to the computation of A_h^1 , we have

$$A_h^3 \leq \lambda e^{-\lambda h} \Phi(0)\Psi(0) \int_0^h \mathbb{E} \left(\mathbf{1}_{\{\tau_x > T_{N_t}\}} v \tilde{f}(t - T_{N_t} + v, x - X_{T_{N_t}}) \right) dv.$$

So,

$$\frac{1}{h} [A_h^1 + A_h^3] \leq \lambda e^{-\lambda h} \Phi(0)\Psi(0) \int_0^h \mathbb{E} \left(\mathbf{1}_{\{\tau_x > T_{N_t}\}} \tilde{f}(t - T_{N_t} + v, x - X_{T_{N_t}}) \right) dv.$$

Using the fact that the application

$$v \mapsto \mathbb{E} \left(\mathbf{1}_{\{\tau_x > T_{N_t}\}} \tilde{f}(t - T_{N_t} + v, x - X_{T_{N_t}}) \right) \text{ is continuous.}$$

yields

$$\lim_{h \rightarrow 0} \frac{1}{h} [A_h^1 + A_h^3] = 0.$$

STEP 2: We now deal with A_h^2 . On the set $\{\tau_x = T_{n+1}\}$, we have:

$$K_x = X_{\tau_x} - x = X_{T_{n+1}} - x = X_{T_n} + [mS_1 + W_{S_1} + Y_1] \circ \Theta_{T_n} - x$$

and

$$L_x = x - X_{\tau_x^-} = x - X_{T_n} - [mS_1 + W_{S_1}] \circ \Theta_{T_n}$$

where Θ is the shift operator. Since $\tilde{X}_{S_1} = [mS_1 + W_{S_1}]$, then

$$K_x = X_{T_n} + [\tilde{X}_{S_1} + Y_1] \circ \Theta_{T_n} - x \text{ and } L_x = x - X_{T_n} - \tilde{X}_{S_1} \circ \Theta_{T_n}.$$

So A_h^2 can be written as following: $A_h^2 =$

$$\sum_{n \geq 0} \mathbb{E} \left(\mathbf{1}_{\tau_x = T_n + S_{n+1} > T_n} \Phi(X_{T_n} + \tilde{X}_{S_{n+1}} + Y_{n+1} - x) \Psi(x - X_{T_n} - \tilde{X}_{S_{n+1}}) \mathbf{1}_{T_n \leq t < T_n + S_{n+1} \leq t+h < T_n + S_{n+1} + S_{n+2}} \right).$$

Strong Markov property applied at T_n leads to $A_h^2 =$

$$\sum_{n \geq 0} \mathbb{E} \left(\mathbf{1}_{\tau_x > T_n} \mathbf{1}_{T_n \leq t} \mathbb{E}^{T_n} (\Phi(X_{T_n} + \tilde{X}_{S_{n+1}} + Y_{n+1} - x) \Psi(x - X_{T_n} - \tilde{X}_{S_{n+1}}) \mathbf{1}_{\tilde{\tau}_{x-X_{T_n}} = S_{n+1}} \mathbf{1}_{t-T_n < S_{n+1} \leq t+h-T_n < S_{n+1} + S_{n+2}}) \right).$$

integrating with respect to S_{n+2} , we have $e^{\lambda h} A_h^2 =$

$$\sum_{n \geq 0} \mathbb{E} \left(\mathbf{1}_{\tau_x > T_n} \mathbf{1}_{T_n \leq t} e^{-\lambda(t-T_n)} \mathbb{E}^{T_n} (e^{\lambda S_{n+1}} \Phi(X_{T_n} + \tilde{X}_{S_{n+1}} + Y_{n+1} - x) \Psi(x - X_{T_n} - \tilde{X}_{S_{n+1}}) \mathbf{1}_{\tilde{\tau}_{x-X_{T_n}} = S_{n+1}} \mathbf{1}_{t-T_n < S_{n+1} \leq t+h-T_n}) \right).$$

We observe that on the set $\{\tilde{\tau}_{x-X_{T_n}} = S_{n+1}\}$, $Y_{n+1} > 0$, $K_x \geq 0$, $L_x \geq 0$ and $K_x + L_x = Y_{n+1}$.

More over:

$$\{\tilde{\tau}_{x-X_{T_n}} = S_{n+1}\} = \left\{ \sup_{s \leq S_{n+1}} \tilde{X}_s < x - X_{T_n}, \tilde{X}_{S_{n+1}} + Y_{n+1} > x - X_{T_n} \right\}.$$

Integrating with respect to (S_{n+1}, Y_{n+1}) implies that

$$\begin{aligned} & \mathbb{E}^{T_n} (e^{\lambda S_{n+1}} \Phi(X_{T_n} + \tilde{X}_{S_{n+1}} + Y_{n+1} - x) \Psi(x - X_{T_n} - \tilde{X}_{S_{n+1}}) \mathbf{1}_{\{\tilde{\tau}_{x-X_{T_n}} = S_{n+1}\}} \mathbf{1}_{t-T_n < S_{n+1} \leq t+h-T_n}) = \\ & \int_0^{+\infty} \int_{t-T_n}^{t+h-T_n} \lambda \mathbb{E}^{T_n} [\Phi(X_{T_n} + \tilde{X}_u + y - x) \Psi(x - X_{T_n} - \tilde{X}_u) \mathbf{1}_{\{\sup_{s \leq u} \tilde{X}_s < x - X_{T_n}, \tilde{X}_u + y > x - X_{T_n}\}}] F_Y(dy) du. \end{aligned}$$

According to Corollary 3.2.1.2 page 147 of [9], $(\sup_{s \leq u} \tilde{X}_s, \tilde{X}_u)$ admits a density

$$\tilde{p}(b, a, t) = \frac{2(2b-a)}{\sqrt{2\pi t^3}} \exp \left[-\frac{(2b-a)^2}{2t} + ma - \frac{m^2}{2}t \right] \mathbf{1}_{b > \max\{0, a\}}.$$

So, A_h^2 is equal to

$$e^{-\lambda h} \sum_{n \geq 0} \mathbb{E} \left[\mathbf{1}_{\tau_x > T_n} \mathbf{1}_{T_n \leq t} e^{-\lambda(t-T_n)} \int_{\mathbb{R}^2} \int_0^{+\infty} \int_{t-T_n}^{t+h-T_n} \lambda \Phi(X_{T_n} + a + y - x) \Psi(x - X_{T_n} - a) \mathbf{1}_{\{b < x - X_{T_n}, a + y > x - X_{T_n}\}} \tilde{p}(b, a, u) \right] F_Y(dy) du db da.$$

Since $e^{-\lambda(t-T_n)} = \mathbb{E}^{T_n}(\mathbf{1}_{T_{n+1}>t})$, and on the event $\{N_t = n\}$, $T_{N_t+1} > t$ a.s, we have

$$A_h^2 = e^{-\lambda h} \mathbb{E} \left[\mathbf{1}_{\tau_x > T_{N_t}} \int_{\mathbb{R}^2} \int_0^{+\infty} \int_{t-T_{N_t}}^{t+h-T_{N_t}} \lambda \Phi(X_{T_{N_t}} + a + y - x) \Psi(x - X_{T_{N_t}} - a) \mathbf{1}_{\{b < x - X_{T_{N_t}}, a + y > x - X_{T_{N_t}}\}} \tilde{p}(b, a, u) \right] F_Y(dy) du db da.$$

We compute the integral with respect to db and it follows

$$A_h^2 = e^{-\lambda h} \mathbb{E} \left[\mathbf{1}_{\tau_x > T_{N_t}} \int_{x-y-X_{T_{N_t}}}^{x-X_{T_{N_t}}} \int_0^{+\infty} \int_{t-T_{N_t}}^{t+h-T_{N_t}} \Phi(X_{T_{N_t}} + a + y - x) \Psi(x - X_{T_{N_t}} - a) \frac{\lambda e^{-\frac{(a-mu)^2}{2u}}}{\sqrt{2\pi u}} F_Y(dy) du da \right] - e^{-\lambda h} \mathbb{E} \left[\mathbf{1}_{\tau_x > T_{N_t}} \int_{x-y-X_{T_{N_t}}}^{x-X_{T_{N_t}}} \int_0^{+\infty} \int_{t-T_{N_t}}^{t+h-T_{N_t}} \Phi(X_{T_{N_t}} + a + y - x) \Psi(x - X_{T_{N_t}} - a) \frac{\lambda e^{-\frac{(a-mu-2x+2X_{T_{N_t}})^2}{2u} + 2m(x-X_{T_{N_t}})}}{\sqrt{2\pi u}} F_Y(dy) du da \right].$$

Change of variables $v = u - (t - T_{N_t})$ and $l = x - X_{T_{N_t}} - a$ yields

$$A_h^2 = e^{-\lambda h} \mathbb{E} \left[\int_0^y \int_0^{+\infty} \int_0^h \mathbf{1}_{\{\tau_x > T_{N_t}\}} \Phi(y-l) \Psi(l) \frac{\lambda e^{-\frac{(x-l-X_{T_{N_t}}-m(t-T_{N_t}+v))^2}{2(t-T_{N_t}+v)}}}{\sqrt{2\pi(t-T_{N_t}+v)}} dv F_Y(dy) dl \right] - e^{-\lambda h} \mathbb{E} \left[\int_0^y \int_0^{+\infty} \int_0^h \mathbf{1}_{\{\tau_x > T_{N_t}\}} \Phi(y-l) \Psi(l) \frac{\lambda e^{-\frac{(x-X_{T_{N_t}}+l+m(t-T_{N_t}+v))^2}{2(t-T_{N_t}+v)} + 2m(x-X_{T_{N_t}})}}}{\sqrt{2\pi(t-T_{N_t}+v)}} dv F_Y(dy) dl \right].$$

Let F_l be the image of F_Y by the map $y \mapsto y - l$. Hence,

$$\lim_{h \rightarrow 0} \frac{1}{h} A_h^2 = \int_0^{+\infty} \int_0^{+\infty} \Phi(k) \Psi(l) \mathbb{E} \left[\mathbf{1}_{\{\tau_x > T_{N_t}\}} \frac{\lambda e^{-\frac{(x-l-X_{T_{N_t}}-m(t-T_{N_t}))^2}{2(t-T_{N_t})}}}{\sqrt{2\pi(t-T_{N_t})}} \right] F_l(dk) dl - \int_0^{+\infty} \int_0^{+\infty} \Phi(k) \Psi(l) \mathbb{E} \left[\mathbf{1}_{\{\tau_x > T_{N_t}\}} \frac{\lambda e^{-\frac{(x-X_{T_{N_t}}+l+m(t-T_{N_t}))^2}{2(t-T_{N_t})} + 2m(x-X_{T_{N_t}})}}}{\sqrt{2\pi(t-T_{N_t})}} \right] F_l(dk) dl$$

since

$$v \mapsto \int_0^{+\infty} \int_0^{+\infty} \Phi(k) \Psi(l) \mathbb{E} \left[\mathbf{1}_{\{\tau_x > T_{N_t}\}} \frac{\lambda e^{-\frac{(x-l-X_{T_{N_t}}-m(t-T_{N_t}+v))^2}{2(t-T_{N_t}+v)}}}{\sqrt{2\pi(t-T_{N_t}+v)}} \right] F_l(dk) dl - \int_0^{+\infty} \int_0^{+\infty} \Phi(k) \Psi(l) \mathbb{E} \left[\mathbf{1}_{\{\tau_x > T_{N_t}\}} \frac{\lambda e^{-\frac{(x-X_{T_{N_t}}-l+l+m(t-T_{N_t}+v))^2}{2(t-T_{N_t}+v)} + 2m(x-X_{T_{N_t}})}}}{\sqrt{2\pi(t-T_{N_t}+v)}} \right] F_l(dk) dl$$

is continuous. $\square$

This proposition concludes the proof of Proposition 5. $\square$

5 Conclusion

Our study relies on the default time of a Lévy process. We have first shown that the distribution function of the default time τ_x belongs to $\mathcal{C}(\mathbb{R}_+^* \times \mathbb{R}_+^*)$ and for any $x \in \mathbb{R}_+^*$, to $\mathcal{C}(\mathbb{R}_+)$. This will be very useful in our future works on default time of a Lévy process where we will use the filtering theory. Secondly, we have obtained an explicit expression to characterize the joint law of the hitting time, overshoot and undershoot of one Lévy process. In this expression, the Gaussian density is of great importance. This law gives a lot of information on the deficit and surplus at default time. In a following paper in progress, we will give a partial differential equation for a Lévy process and its running maximum.

6 Appendix

Lemma 1. *Let $\beta > -1$, then for $0 < t \leq 1$,*

$$\mathbb{E} \left(\mathbf{1}_{\{N_t \geq 2\}} (t - T_{N_t})^\beta \right) \leq \left(\sum_{n=1}^{\infty} \frac{\lambda^n e^t}{(n-1)!} B(n, \beta + 1) \right) t^{2+\beta}$$

where $B(n, \beta + 1) = \int_0^1 (1-u)^\beta u^{n-1} du$.

Proof. According to the values of the process N_t , we have

$$\begin{aligned} \mathbb{E} \left(\mathbf{1}_{\{N_t \geq 2\}} (t - T_{N_t})^\beta \right) &= \sum_{n \geq 2} \mathbb{E} \left((t - T_n)^\beta \mathbf{1}_{T_n \leq t < T_n + S_{n+1}} \right) \\ &= \sum_{n \geq 2} \mathbb{E} \left(e^{-\lambda(t-T_n)} (t - T_n)^\beta \right). \end{aligned}$$

Since T_n follows a Gamma law of parameters n and λ , hence

$$\mathbb{E} \left(\mathbf{1}_{\{N_t \geq 2\}} (t - T_{N_t})^\beta \right) = \sum_{n=2}^{\infty} \frac{\lambda^n e^{-\lambda t}}{(n-1)!} t^{n+\beta} B(n, \beta + 1).$$

□

Lemma 2. *For any $0 < t_1 < t_2$,*

$$\sup_{t_1 \leq t \leq t_2} \mathbb{E} \left(\mathbf{1}_{\tau_x > T_{N_t}} \tilde{f}(t - T_{N_t}, x - X_{T_{N_t}})^p \right) < +\infty. \quad (15)$$

and for instance, as a consequence, the family

$$\left(\mathbf{1}_{\tau_x > T_{N_t}} \tilde{f}(t - T_{N_t}, x - X_{T_{N_t}}), \quad t \in [t_1, t_2], \quad x > 0 \right)$$

is uniformly integrable.

Proof. Refer to Lemma 3.1 in [5], if G is a Gaussian random variable $\mathcal{N}(0, 1)$, $\mu > 0$, $u > 0$, $\sigma \in \mathbb{R}^+$ and $p \geq 1$, then

$$\mathbb{E}[(\tilde{f}(u, \mu + \sigma G))^p 1_{\mu + \sigma G > 0}] = \frac{1}{\sqrt{(2\pi)^p}} \frac{u^{\frac{1-2p}{2}} e^{-\frac{p(\mu - mu)^2}{2(p\sigma^2 + u)}}}{(p\sigma^2 + u)^{\frac{p+1}{2}}} \mathbb{E} \left[\left(\sigma G + \sqrt{\frac{u}{p\sigma^2 + u}} (\mu - mu) + m \sqrt{u(p\sigma^2 + u)} \right)_+^p \right]$$

Using the inequality $(a + b + c)^p \leq 3^{p-1}(a^p + b^p + c^p)$ where a, b, c are positive numbers, it follows that

$$\mathbb{E}[\tilde{f}(u, \mu + \sigma G)^p 1_{\mu + \sigma G > 0}] \leq \frac{3^{p-1}}{\sqrt{(2\pi)^p}} \frac{u^{\frac{1-2p}{2}} e^{-\frac{p(\mu - mu)^2}{2(p\sigma^2 + u)}}}{(p\sigma^2 + u)^{\frac{p+1}{2}}} \left(\sigma^p \mathbb{E}(|G|^p) + \left(\frac{u}{p\sigma^2 + u}\right)^{\frac{p}{2}} (\mu - mu)^p + m^p (u(p\sigma^2 + u))^{\frac{p}{2}} \right).$$

Therefore

$$\begin{aligned} \mathbb{E}[\tilde{f}(u, \mu + \sigma G)^p 1_{\mu + \sigma G > 0}] &\leq \frac{3^{p-1}}{\sqrt{2\pi}^p} \left(\frac{u^{\frac{1-2p}{2}}}{(p\sigma^2 + u)^{\frac{p+1}{2}}} \sigma^p \mathbb{E}(|G|^p) + \frac{u^{\frac{1-p}{2}}}{(p\sigma^2 + u)^{\frac{p+1}{2}}} c_p + |m|^p \frac{u^{\frac{1-p}{2}}}{(p\sigma^2 + u)^{\frac{1}{2}}} \right) \\ &\leq \frac{3^{p-1}}{\sqrt{2\pi}^p} \left(\frac{u^{\frac{1-2p}{2}}}{(p\sigma^2 + u)^{\frac{p+1}{2}}} \sigma^p \mathbb{E}(|G|^p) + \frac{c_p}{u^{\frac{p-1}{2}} (p\sigma^2 + u)^{\frac{p+1}{2}}} + \frac{|m|^p}{u^{\frac{p-1}{2}} (p\sigma^2 + u)^{\frac{1}{2}}} \right). \end{aligned}$$

Using the independence between the Brownian motion and the Poisson process, we can apply this inequality to $\sigma = \sqrt{T_{N_t}}$, $u = t - T_{N_t}$, $p\sigma^2 + u = (p-1)T_{N_t} + t \geq t > t_1$:

$$\begin{aligned} &| \mathbb{E}(1_{\tau_x > T_{N_t}} \tilde{f}(t - T_{N_t}, x - X_{N_t})^p) | \leq \\ &\frac{3^{p-1}}{\sqrt{2\pi}^p} \mathbb{E} \left(\frac{T_{N_t}^{\frac{p}{2}} \mathbb{E}(|G|^p)}{(t - T_{N_t})^{\frac{2p-1}{2}} t^{\frac{p+1}{2}}} + \frac{c_p}{(t - T_{N_t})^{\frac{p-1}{2}} t^{\frac{2p+1}{2}}} + \frac{|m|^p}{\sqrt{t} (t - T_{N_t})^{\frac{p-1}{2}}} \right). \end{aligned}$$

We use the following for $\alpha = 0$ or $p/2$, and $\beta = (2p-1)/2$ or $(p-1)/2$:

$$\begin{aligned} \mathbb{E}\left(\frac{T_{N_t}^\alpha}{(t - T_{N_t})^\beta}\right) &= \frac{1}{t^\beta} + \sum_{n \geq 1} \mathbb{E}\left[\frac{T_n^\alpha}{(t - T_n)^\beta} 1_{T_n < t < T_{n+1}}\right] \\ &= \frac{1}{t^\beta} + \sum_{n \geq 1} \int_0^t \frac{u^\alpha}{(t-u)^\beta} \frac{(\lambda u)^{n-1}}{(n-1)!} \lambda e^{-\lambda u} \int_{t-u}^{+\infty} \lambda e^{-\lambda v} dv du \\ &= \frac{1}{t^\beta} + e^{-\lambda(t)} \sum_{n \geq 1} \frac{\lambda^n}{(n-1)!} \int_0^t \frac{u^{n+\alpha-1}}{(t-u)^\beta} du \\ &= \frac{1}{t^\beta} + e^{-\lambda(t)} \sum_{n \geq 1} \frac{[\lambda(t)]^n}{(n-1)!} B(\alpha + n; 1 - \beta) (t)^{\alpha-\beta} \end{aligned}$$

where $B(n, \beta + 1) = \int_0^1 (1-u)^\beta u^{n-1} du$. Since $t_2 \geq t \geq t_1 > 0$, we conclude. $\square$

Lemma 3. For any numbers u, v, y, z and a , the equality

$$\frac{1}{u}(a-z)^2 + \frac{1}{v}(a-y)^2 = \frac{v+u}{uv} \left[a - \frac{vz+uy}{v+u} \right]^2 + \frac{1}{u+v}(z-y)^2$$

holds.

Proof. We develop both squared

$$\frac{1}{u}(a-z)^2 + \frac{1}{v}(a-y)^2 = \left(\frac{1}{u} + \frac{1}{v}\right)a^2 - 2a\left(\frac{z}{u} + \frac{y}{v}\right) + \frac{z^2}{u} + \frac{y^2}{v}.$$

We have the first square

$$\begin{aligned} \frac{1}{u}(a-z)^2 + \frac{1}{v}(a-y)^2 &= \left(\frac{1}{u} + \frac{1}{v}\right) \left[a^2 - 2a\left(\frac{z}{u} + \frac{y}{v}\right)\frac{uv}{v+u} + \left[\left(\frac{z}{u} + \frac{y}{v}\right)\frac{uv}{v+u}\right]^2 \right] \\ &\quad + \frac{z^2}{u} + \frac{y^2}{v} - \left[\frac{z^2}{u^2} + 2\frac{yz}{uv} + \frac{y^2}{v^2} \right] \frac{uv}{u+v}. \end{aligned}$$

We order

$$\begin{aligned} \frac{1}{u}(a-z)^2 + \frac{1}{v}(a-y)^2 &= \left(\frac{1}{u} + \frac{1}{v}\right) \left[a^2 - 2a\left(\frac{z}{u} + \frac{y}{v}\right)\frac{uv}{v+u} + \left[\left(\frac{z}{u} + \frac{y}{v}\right)\frac{uv}{v+u}\right]^2 \right] \\ &\quad + z^2 \frac{1}{u} \left[1 - \frac{v}{u+v} \right] + y^2 \frac{1}{v} \left[1 - \frac{u}{u+v} \right] - 2\frac{yz}{uv} \frac{uv}{u+v}. \end{aligned}$$

which concludes the proof. $\square$

Lemma 4. Let be $(\tilde{X}_u, u \geq 0)$ be a Brownian motion with drift $m \in \mathbb{R}$. So, we have

$$\mathbb{E} \left(\mathbf{1}_{\{\tilde{X}_u^* < c\}} | \tilde{X}_u \right) = \mathbf{1}_{\{\tilde{X}_u < c\}} \left[1 - \exp \left[-\frac{2c^2}{u} + \frac{2c}{u} \tilde{X}_u \right] \right]$$

for all real number $u > 0$.

Proof. Refer to Corollary 3.2.1.2 page 147 of [9], $(\tilde{X}_t^*, \tilde{X}_t)$ admits a density

$$\tilde{p}(b, a, t) = \frac{2(2b-a)}{\sqrt{2\pi t^3}} \exp \left[-\frac{(2b-a)^2}{2t} + ma - \frac{m^2}{2}t \right] \mathbf{1}_{b > \max\{0, a\}}$$

So, the conditional law of $\tilde{X}_u^*$ given $\tilde{X}_u = a$ has the density

$$f_{\tilde{X}^* | \tilde{X}=a}(b|a) = \frac{2(2b-a)}{u} \exp \left[-\frac{(2b-a)^2 - a^2}{2u} \right] \mathbf{1}_{b > \max\{0, a\}}.$$

Thus

$$\begin{aligned} \mathbb{E} \left(\mathbf{1}_{\{\tilde{X}_u^* < c\}} | \tilde{X}_u = a \right) &= \{ \mathbf{1}_{a < c} \} \int_{\max\{a, 0\}}^c \frac{2(2b-a)}{u} \exp \left[-\frac{(2b-a)^2 - a^2}{2u} \right] \mathbf{1}_{b > \max\{0, a\}} db \\ &= \mathbf{1}_{\{a < c\}} \left[1 - \exp \left[-\frac{2c^2}{u} + \frac{2c}{u}a \right] \right]. \end{aligned}$$

$\square$

Acknowledgments We wish to thank Monique Pontier for her carefully reading this paper and making useful suggestions.

References

- [1] J. Bertoin (1996), *Lévy Processes*. Cambridge Univ. Press. MR 1406504.
- [2] V. Bernyk, R. C. Dalang, G. Peskir, *The law of the supremum of stable Lévy processes with no negative jumps* Ann. Proba. Vol 36, Number 5, 2008, pp. 177-1789.
- [3] A. A. Borovkov, *On the first passage time for one class of processes with independent increments*, Theory of Probability & Its Applications, Vol. 10, N 32, pages 331–334, 1964, SIAM
- [4] R. Cont and P. Tankov, *Financial Modeling with jump Processes*, Chapman & Hall/ CRC, Financial Mathematics Series, 2004.
- [5] L. Coutin and D. Dorobantu, *First passage law for some Lévy processes with compound Poisson: Existence of a density*, Bernoulli, vol 17, N 4, 2011, pp 1127–1135.
- [6] R. A. Doney, *Hitting probabilities for spectrally positive lévy processes*, J. London Math. Soc. (2) 44. 1991, pp. 556-576.
- [7] R. A. Doney and A. E. Kyprianou, *Overshoot and undershoot of Lévy processes*, Ann. Appl. Probab 2006, Vol. 16, No. 1, pp 91-106.
- [8] R. A. Doney, *A note on the supremum of a stable Lévy process*, stochastics, (2008), 80, 151-155. MR-2402160.
- [9] M. Jeanblanc, M. Yor, M. Chesney, *Mathematical methods for financial markets*, 2009, Springer.
- [10] I. Karatzas and S.E Shreve, *Brownian motion and stochastic calculus*, Vol 113, Springer Science & Business Media, 2012.
- [11] S. G. Kou and H. Wang, *First passage time of jump diffusion process*, Adv. Appl. Prob. 35, 2003, pp. 504-531.
- [12] Kuznetsov Alexey, Kyprianou Andreas E, Pardo Juan Carlos, Watson Alexander R., *The hitting time of zero for a stable process*, Electron. J. Probab, vol 19, Number 30, pp 1–26, 2014.
- [13] A. E. Kyprianou, *Fluctuations of Lévy Processes with Applications: Introductory Lectures*, Springer Science & Business Media, 2014.
- [14] G. Peskir, *The law of the hitting times to points by a stable Lévy process with no-negative jumps*, Research Report No. 15, Probability and statistics Group school of Mathematics, The university of Manchester, 2007.
- [15] Pogány, Tibor K and Nadarajah, Saralees, *On the result of Doney*, Electronic communications in probability, vol. 20, pp. 1-4, 2015

- [16] B. Roynette, P. Vallois, A. Volpi, *Asymptotic Behavior of the hitting time, overshoot and undershoot for some Lévy processes* *ESAIM PS Vol 12, 2008* pp 58-98.
- [17] V. M. Zolotarev, *The first passage time of a level and the behavior at infinity for a class of processes with independent increments*, *Theory of Probability & Its Applications, Vol. 9, 1964*, pp. 653-662, SIAM.